

Preoperative Instructions for Blood Thinners

Please **STOP ALL BLOOD THINNERS 2 WEEKS before your procedure.**

Some of these include:

ASPIRIN	DAYPRO	NAPPRELAN
ADVIL	DOLOBID	NUPRIN
AGGRENEX	ECOTRIN	NAFLON
ALEVE	FELDENE	ORUDIS
ALKA SELTZER	IBUPROFEN	PERSANTINE
ANSAID	INDOCIN	PLETAL
ANAPROX	INNOHEP	PONSTEL
BEXTRA	LODINE	RELAFEN
BUFFERIN	MOBIC	TICLID
CATAFLAM	MOTRIN	TELECTIN
CELEBREX	NAPROSYN	TORADEL
CLINORIL	NAPROXEN	VOLTAREN
DICLOFENAC	MELOXIAM	

**ALL VITAMINS INCLUDING:
MULTI-VITAMINS, VITAMIN E
GLUCOSAMINE
ALL HOMEOPATHIC REMEDIES
ALL HERBS**

**ARGATROBAN
BIVALIRUDIN (ANGIOMAX)
LEPIRUDIN (REFLUDAN)**

EMBRIL - HOLD 2 WEEK PRIOR

Please **STOP THE FOLLOWING 5 DAYS before your procedure:**

COUMADIN PLAVIX WARFARIN

IF YOU HAVE ANY QUESTIONS, PLEASE CALL THE OFFICE AT 215-257-3066.

THANK YOU FOR YOUR COOPERATION.

Patient Signature: _____ Date: _____

Patient Name (Printed): _____

Rev 6/09 tk

711 Lawn Avenue
Building 2
Sellersville, PA 18960

2031 N. Broad Street
Suite 141
Lansdale, PA 19446

1021 Park Avenue
Suite 30
Quakertown, PA 18951

T 215.257.3066
215.257.1050
F 215.257.1634
215.257.3026